CLAIMS

1. A compound of the formula (I):

$$R^{1}$$
 $A-N$
 R^{3}
 E
 $N-N$
 H
 (I)

or a salt, solvate, tautomer or N-oxide thereof;

5

wherein A is a saturated hydrocarbon linker group containing from 1 to 7 carbon atoms, the linker group having a maximum chain length of 5 atoms extending between R^1 and NR^2R^3 and a maximum chain length of 4 atoms extending between E and NR^2R^3 , wherein one of the carbon atoms in the linker group may optionally be replaced by an oxygen or nitrogen atom; and wherein the carbon atoms of the linker group A may optionally bear one or more substituents selected from oxo, fluorine and hydroxy, provided that the hydroxy group when present is not located at a carbon atom α with respect to the NR^2R^3 group and provided that the oxo group when present is located at a carbon atom α with respect to the NR^2R^3 group;

15

10

E is a monocyclic or bicyclic carbocyclic or heterocyclic group wherein E is unsubstituted or has up to 4 substituents R^8 selected from hydroxy, oxo (when E is non-aromatic), chlorine, bromine, trifluoromethyl, cyano, C_{1-4} hydrocarbyloxy and C_{1-4} hydrocarbyl optionally substituted by C_{1-2} alkoxy or hydroxyl;

20

 R^1 is an aryl or heteroaryl group which is unsubstituted or bears one or more substituents selected from hydroxy; C_{1-4} acyloxy; fluorine; chlorine; bromine; trifluoromethyl; cyano; CONH₂; nitro; C_{1-4} hydrocarbyloxy and C_{1-4} hydrocarbyl each optionally substituted by C_{1-2} alkoxy, carboxy or hydroxy; C_{1-4}

acylamino; benzoylamino; pyrrolidinocarbonyl; piperidinocarbonyl; morpholinocarbonyl; piperazinocarbonyl; five and six membered heteroaryl and heteroaryloxy groups containing one or two heteroatoms selected from N, O and S; phenyl; phenyl-C₁₋₄ alkyl; phenyl-C₁₋₄ alkoxy; heteroaryl-C₁₋₄ alkyl; heteroaryl-C₁₋₄ alkoxy and phenoxy, wherein the heteroaryl, heteroaryloxy, phenyl, phenyl-C₁₋₄ alkyl, phenyl-C₁₋₄ alkoxy, heteroaryl-C₁₋₄ alkyl, heteroaryl-C₁₋₄ alkoxy and phenoxy groups are each optionally substituted with 1, 2 or 3 substituents selected from C₁₋₂ acyloxy, fluorine, chlorine, bromine, trifluoromethyl, cyano, CONH₂, C₁₋₂ hydrocarbyloxy and C₁₋₂ hydrocarbyl each optionally substituted by methoxy or hydroxyl;

 R^2 and R^3 are independently selected from hydrogen, $C_{1.4}$ hydrocarbyl and $C_{1.4}$ acyl wherein the hydrocarbyl and acyl moieties are optionally substituted by one or more substituents selected from fluorine, hydroxy, amino, methylamino, dimethylamino and methoxy;

or R² and R³ together with the nitrogen atom to which they are attached form a cyclic group selected from an imidazole group and a saturated monocyclic heterocyclic group having 4-7 ring members and optionally containing a second heteroatom ring member selected from O and N;

or one of R² and R³ together with the nitrogen atom to which they are attached and one or more atoms from the linker group A form a saturated monocyclic heterocyclic group having 4-7 ring members and optionally containing a second heteroatom ring member selected from O and N;

or NR²R³ and the carbon atom of linker group A to which it is attached together form a cyano group;

 R^4 is selected from hydrogen, halogen, C_{1-5} saturated hydrocarbyl, C_{1-5} saturated hydrocarbyloxy, cyano, and CF_3 ; and

R⁵ is selected from hydrogen, halogen, C₁₋₅ saturated hydrocarbyl, C₁₋₅ saturated hydrocarbyloxy, cyano, CONH₂, CONHR⁹, CF₃, NH₂, NHCOR⁹ or NHCONHR⁹;

. 10

5

15

20

 R^9 is a group R^{9a} or $(CH_2)R^{9a}$, wherein R^{9a} is a monocyclic or bicyclic group which may be carbocyclic or heterocyclic;

the carbocyclic group or heterocyclic group R^{9a} being optionally substituted by one or more substituents selected from halogen, hydroxy, trifluoromethyl, cyano, nitro, carboxy, amino, mono- or di- C_{1-4} hydrocarbylamino; a group R^a - R^b wherein R^a is a bond, O, CO, $X^1C(X^2)$, $C(X^2)X^1$, $X^1C(X^2)X^1$, S, SO, SO₂, NR^c , SO₂ NR^c or NR^c SO₂; and R^b is selected from hydrogen, heterocyclic groups having from 3 to 12 ring members, and a C_{1-8} hydrocarbyl group optionally substituted by one or more substituents selected from hydroxy, oxo, halogen, cyano, nitro, carboxy, amino, mono- or di- C_{1-4} hydrocarbylamino, carbocyclic and heterocyclic groups having from 3 to 12 ring members and wherein one or more carbon atoms of the C_{1-8} hydrocarbyl group may optionally be replaced by O, S, SO, SO₂, NR^c , $X^1C(X^2)$, $C(X^2)X^1$ or $X^1C(X^2)X^1$;

15

10

5

R^c is selected from hydrogen and C₁₋₄ hydrocarbyl; and
X¹ is O, S or NR^c and X² is =O, =S or =NR^c;
but excluding the compound (2R, 5S)-1-benzyl-4-(R)-1-(3-[1-(tert-butyl)-1,1-dimethylsilyl]oxyphenyl)-1-[4-(1H-pyrazol-4-yl)phenyl]methyl-2,5-dimethylhexahydropyrazine.

20 2. A compound according to claim 1 of the formula (Ia):

$$R^{1}$$
 $A-N$
 R^{3}
 E
 R^{4}
 $N-N$
 H
(Ia)

or a salt, solvate, tautomer or N-oxide thereof;

wherein A is a saturated hydrocarbon linker group containing from 1 to 7 carbon atoms, the linker group having a maximum chain length of 5 atoms extending between R^1 and NR^2R^3 and a maximum chain length of 4 atoms extending between E and NR^2R^3 , wherein one of the carbon atoms in the linker group may optionally be replaced by an oxygen or nitrogen atom; and wherein the carbon atoms of the linker group A may optionally bear one or more substituents selected from oxo, fluorine and hydroxy, provided that the hydroxy group when present is not located at a carbon atom α with respect to the NR^2R^3 group and provided that the oxo group when present is located at a carbon atom α with respect to the NR^2R^3 group;

E is a monocyclic or bicyclic carbocyclic or heterocyclic group wherein E is unsubstituted or has up to 4 substituents R⁸ as defined in claim 1;

R¹ is an aryl or heteroaryl group which is unsubstituted or substituted as defined in claim 1

 R^2 and R^3 are independently selected from hydrogen, C_{1-4} hydrocarbyl and C_{1-4} acyl;

or R² and R³ together with the nitrogen atom to which they are attached form a saturated monocyclic heterocyclic group having 4-7 ring members and optionally containing a second heteroatom ring member selected from O and N;

or one of R² and R³ together with the nitrogen atom to which they are attached and one or more atoms from the linker group A form a saturated monocyclic heterocyclic group having 4-7 ring members and optionally containing a second heteroatom ring member selected from O and N;

or NR²R³ and the carbon atom of linker group A to which it is attached together form a cyano group;

R⁴ is selected from hydrogen, halogen, C₁₋₅ saturated hydrocarbyl, cyano and CF₃; and

R⁵ is selected from hydrogen, halogen, C₁₋₅ saturated hydrocarbyl, cyano, CONH₂, CONHR⁹, CF₃, NH₂, NHCOR⁹ or NHCONHR⁹;

10

5

15

20

 R^9 is phenyl or benzyl each optionally substituted by one or more substituents selected from halogen, hydroxy, trifluoromethyl, cyano, nitro, carboxy, amino, mono- or di- C_{1-4} hydrocarbylamino; a group R^a - R^b wherein R^a is a bond, O, CO, $X^1C(X^2)$, $C(X^2)X^1$, $X^1C(X^2)X^1$, S, SO, SO₂, NR^c , SO₂ NR^c or NR^c SO₂; and R^b is selected from hydrogen, heterocyclic groups having from 3 to 12 ring members, and a C_{1-8} hydrocarbyl group optionally substituted by one or more substituents selected from hydroxy, oxo, halogen, cyano, nitro, carboxy, amino, mono- or di- C_{1-4} hydrocarbylamino, carbocyclic and heterocyclic groups having from 3 to 12 ring members and wherein one or more carbon atoms of the C_{1-8} hydrocarbyl group may optionally be replaced by O, S, SO, SO₂, NR^c , $X^1C(X^2)$, $C(X^2)X^1$ or $X^1C(X^2)X^1$;

5

10

15

R^c is selected from hydrogen and C₁₋₄ hydrocarbyl; and
X¹ is O, S or NR^c and X² is =O, =S or =NR^c;
but excluding the compound (2R, 5S)-1-benzyl-4-(R)-1-(3-[1-(tert-butyl)-1,1-

dimethylsilyl]oxyphenyl)-1-[4-(1H-pyrazol-4-yl)phenyl]methyl-2,5-dimethylhexahydropyrazine.

- A compound according to claim 1 or claim 2 wherein A is a saturated hydrocarbon linker group containing from 1 to 7 carbon atoms, the linker group having a maximum chain length of 5 atoms extending between R¹ and NR²R³ and a maximum chain length of 4 atoms extending between E and NR²R³, wherein one of the carbon atoms in the linker group may optionally be replaced by an oxygen or nitrogen atom; and wherein the carbon atoms of the linker group A may optionally bear one or more substituents selected from fluorine and hydroxy, provided that the hydroxy group when present is not located at a carbon atom α with respect to the NR²R³ group; and R⁵ is selected from selected from hydrogen, halogen, C₁₋₅ saturated hydrocarbyl, cyano, CONH₂, CF₃, NH₂, NHCOR⁹ and NHCONHR⁹.
 - 4. A compound according to any one of claims 1 to 3 wherein the linker group A has a maximum chain length of 3 atoms extending between R¹ and NR²R³.

- 5. A compound according to claim 4 wherein the linker group A has a maximum chain length of 2 atoms extending between R¹ and NR²R³.
- 6. A compound according to any one of claims 1 to 5 wherein the linker group A has a maximum chain length of 3 atoms extending between E and NR²R³.
- A compound according to claim 6 wherein the linker group A has a chain length of 2 or 3 atoms extending between R¹ and NR²R³ and a chain length of 2 or 3 atoms extending between E and NR²R³.
- 8. A compound according to any one of the preceding claims wherein the linker group atom linked directly to the group E is a carbon atom and the linker group A has an all-carbon skeleton.
- 9. A compound according to any one of claims 1 to 7 wherein the portion R¹-A-NR²R³ of the compound is represented by the formula R¹-(G)_k-(CH₂)_m-W-O_b-(CH₂)_n-(CR⁶R³)_p-NR²R³ wherein G is NH, NMe or O; W is attached to the group E and is selected from (CH₂)_j-CR²0, (CH₂)_j-N and (NH)_j-CH; b is 0 or 1, j is 0 or 1, k is 0 or 1, m is 0 or 1, n is 0, 1, 2, or 3 and p is 0 or 1; the sum of b and k is 0 or 1; the sum of j, k, m, n and p does not exceed 4; R⁶ and R³ are the same or different and are selected from methyl and ethyl, or CR⁶R³ forms a cyclopropyl group; and R²0 is selected from hydrogen, methyl, hydroxy and fluorine.
- 10. A compound according to any one of claims 1 to 7 wherein the moiety R¹-A
 NR²R³ is represented by the formula R¹-(G)_k-(CH₂)_m-X-(CH₂)_n-(CR⁶R⁷)_p-NR²R³

 wherein G is NH, NMe or O; X is attached to the group E and is selected from

 (CH₂)_j-CH, (CH₂)_j-N and (NH)_j-CH; j is 0 or 1, k is 0 or 1, m is 0 or 1, n is 0, 1, 2,

 or 3 and p is 0 or 1, and the sum of j, k, m, n and p does not exceed 4; and R⁶ and

 R⁷ are the same or different and are selected from methyl and ethyl, or CR⁶R⁷

 forms a cyclopropyl group.
 - 11. A compound according to claim 10 wherein (i) k is 0, m is 0 or 1, n is 0, 1, 2 or 3 and p is 0; or (ii) k is 0, m is 0 or 1, n is 0, 1 or 2 and p is 1.

- 12. A compound according to claim 10 wherein (i) X is $(CH_2)_j$ -CH, k is 1, m is 0, n is 0, 1,2 or 3 and p is 0; or (ii) X is $(CH_2)_j$ -CH, k is 1, m is 0, n is 0, 1 or 2 and p is 1.
- 13. A compound according to claim 10 or claim 12 wherein (i) j is 0; or (ii) j is 1; or (iii) CR⁶R⁷ is C(CH₃)₂.
- A compound according to claim 10 wherein the portion R¹-A-NR²R³ of the compound is represented by the formula R¹-X-(CH₂)_n-NR²R³ where X is attached to the group E and is a group CH, and n is 2.
 - 15. A compound according to claim 1 or claim 2 wherein R¹-A(E)-NR²R³ is selected from the groups A1 to A11 below:

$R^1 \longrightarrow R^2$ $N \searrow R^3$	R^{1} \uparrow R^{2} \downarrow R^{3}	$R^1 \longrightarrow N^{R^2}$ $E R^3$
E A1	A2	A3
R ¹ Me Me R ² N R ² A4	R^1 Me Me R^2 R^3 $A5$	R^1 R^2 R^3 R^3 R^3 R^4 R^3 R^4 R^3 R^4
R ¹ N R ² R ³ A7	P ¹ OH R ² N R ³ R ³ A8	R ¹ C N A9
R^1 R^2 R^3 R^3 R^3 R^3	R ¹ All	

- 16. A compound according to any one of the preceding claims wherein E is a monocyclic group.
- 17. A compound according to any one of the preceding claims wherein E is an aryl or heteroaryl group each of which is unsubstituted or substituted by up to 4 substituents R⁸ as defined in claim 1.
- 18. A compound according to claim 17 selected from optionally substituted phenyl, thiophene, furan, pyrimidine and pyridine groups, each of which is unsubstituted or substituted by up to 4 substituents R⁸ as defined in claim 1.
- 19. A compound according to claim 18 wherein E is a phenyl group which is unsubstituted or substituted by up to 4 substituents R⁸ as defined in claim 1.

- 20. A compound according to any one of the preceding claims wherein the group A and the pyrazole group are attached to the group E in a *meta* or *para* relative orientation; i.e. A and the pyrazole group are not attached to adjacent ring members of the group E.
- 1.3 A compound according to claim 20 wherein E is selected from 1,4-phenylene, 1,3-phenylene, 2,5-pyridylene and 2,4-pyridylene, 1,4-piperazinyl, and 1,4-piperazonyl, each of which is unsubstituted or substituted by up to 4 substituents R⁸ as defined in claim 1.
- 22. A compound according to any one of the preceding claims wherein E has 0-3 substituents.
 - 23. A compound according to claim 22 wherein E has 0-2 substituents
 - 24. A compound according to claim 23 wherein E has 0 or 1 substituent.
 - 25. A compound according to claim 24 wherein E is unsubstituted.

A compound according to any one of the preceding claims wherein the group E is 26. an aryl or heteroaryl group having five or six members and containing up to three heteroatoms selected from O, N and S, the group E being represented by the formula:

5

where * denotes the point of attachment to the pyrazole group, and "a" denotes the attachment of the group A;

r is 0, 1 or 2;

U is selected from N and CR^{12a}; and

10

V is selected from N and CR^{12b}; where R^{12a} and R^{12b} are the same or different and each is hydrogen or a substituent containing up to ten atoms selected from C, N, O, F, Cl and S provided that the total number of non-hydrogen atoms present in R^{12a} and R^{12b} together does not exceed ten;

or R^{12a} and R^{12b} together with the carbon atoms to which they are attached form an unsubstituted five or six membered saturated or unsaturated ring containing up to two heteroatoms selected from O and N; and

15

R¹⁰ is selected from halogen, hydroxy, trifluoromethyl, cyano, nitro, carboxy, amino, mono- or di-C₁₋₄ hydrocarbylamino, carbocyclic and heterocyclic groups having from 3 to 12 ring members; a group Ra-Rb wherein Ra is a bond, O, CO, X¹C(X²), C(X²)X¹, X¹C(X²)X¹, S, SO, SO₂, NR^c, SO₂NR^c or NR^cSO₂; and R^b is selected from hydrogen, carbocyclic and heterocyclic groups having from 3 to 12 ring members, and a C₁₋₈ hydrocarbyl group optionally substituted by one or more substituents selected from hydroxy, oxo, halogen, cyano, nitro, carboxy, amino, mono- or di-C₁₋₄ hydrocarbylamino, carbocyclic and heterocyclic groups having

20

from 3 to 12 ring members and wherein one or more carbon atoms of the C₁₋₈

hydrocarbyl group may optionally be replaced by O, S, SO, SO₂, NR^c, $X^1C(X^2)$, $C(X^2)X^1$ or $X^1C(X^2)X^1$;

 R^{c} is selected from hydrogen and C_{1-4} hydrocarbyl; and X^{1} is O, S or NR^{c} and X^{2} is =O, =S or = NR^{c} .

5 27. A compound according to claim 26 wherein E is represented by the formula:

where P, Q and T are the same or different and are selected from N, CH and NCR¹⁰, provided that the group A is attached to a carbon atom.

28. A compound according to claim 27 wherein the group E is selected from groups
10 B1 to B13 in the Table below:

* B1	B2	B3	a N N * B4
* B5	a * B6	* B7	B8

MeO * B9	R ¹³ * B10	R ¹³ * B11	B12
B13			

29. A compound according to claim 19 having the formula (II):

$$R^{1}$$
 R^{2}
 R^{3}
 R^{4}
 R^{4}
 R^{5}
 R^{5}
 R^{1}
 R^{2}
 R^{3}
 R^{4}
 R^{5}
 R^{1}
 R^{2}
 R^{3}
 R^{4}
 R^{5}
 R^{1}
 R^{2}
 R^{3}
 R^{4}
 R^{5}
 R^{5}
 R^{1}
 R^{5}
 R^{5}
 R^{5}
 R^{7}
 R^{7}

wherein the group A is attached to the *meta* or *para* position of the benzene ring and q is 0-4; R^8 is hydroxy; halogen; trifluoromethyl; cyano; C_{1-4} hydrocarbyloxy optionally substituted by C_{1-2} alkoxy or hydroxy; and C_{1-4} hydrocarbyl optionally substituted by C_{1-2} alkoxy or hydroxy.

30. A compound according to claim 29 wherein q is 0, 1 or 2

5

31. A compound according to claim 30 wherein q is 0 or 1.

- 32. A compound according to claim 31 wherein q is 0.
- 33. A compound according to claim 19 having the formula (III):

$$R^{1}$$
 A'
 N
 R^{3}
 R^{3}
 R^{5}
 $N-N$
 H
(III)

where A' is the residue of the group A and R¹ to R⁵ are as defined in any one of the preceding claims.

34. A compound according to claim 19 having the formula (IV):

5

$$R^{1}$$
 R^{20}
 R^{2}
 R^{3}
 R^{4}
 R^{5}
 R^{5}
 R^{5}
 R^{5}
 R^{5}
 R^{7}
 R^{1}
 R^{2}
 R^{2}
 R^{3}

wherein z is 0, 1 or 2, R^{20} is selected from hydrogen, methyl, hydroxy and fluorine, provided that when z is 0, R^{20} is other than hydroxy.

10 35. A compound according to claim 19 having the formula (V):

wherein R³ is optionally selected from hydrogen and C₁₋₄ hydrocarbyl.

- 36. A compound according to claim 35 wherein \mathbb{R}^3 is selected from hydrogen and \mathbb{C}_{1-4} hydrocarbyl.
- 5 37. A compound according to claim 29 wherein R¹ is phenyl.
 - 38. A compound according to any one of the preceding claims wherein R¹ is selected from phenyl, naphthyl, thienyl, furan, pyrimidine and pyridine, each optionally substituted as defined in claim 1.
- 39. A compound according to any one of the preceding claims wherein R¹ is

 10 unsubstituted or is substituted by up to 5 substituents selected from hydroxy; C₁₋₄

 acyloxy; fluorine; chlorine; bromine; trifluoromethyl; cyano; C₁₋₄ hydrocarbyloxy

 and C₁₋₄ hydrocarbyl optionally substituted by C₁₋₂ alkoxy or hydroxy; and five

 membered heteroaryl groups containing one or two heteroatoms selected from N,

 O and S, the heteroaryl groups being optionally substituted by one or more C₁₋₄

 alkyl substituents.
 - 40. A compound according to claim 23 wherein R¹ is unsubstituted or is substituted by up to 5 substituents selected from hydroxy, C₁₋₄ acyloxy, fluorine, chlorine, bromine, trifluoromethyl, cyano, C₁₋₄ hydrocarbyloxy and C₁₋₄ hydrocarbyl optionally substituted by C₁₋₂ alkoxy or hydroxy.

- 41. A compound according to claim 23 or 24 wherein the group R¹ has one or two substituents selected from fluorine, chlorine, trifluoromethyl, methyl and methoxy.
- 42. A compound according to claim 25 wherein R¹ is a mono-chlorophenyl or dichlorophenyl group.

- 43. A compound according to any one of the preceding claims wherein R⁴ is selected from hydrogen and methyl.
- 44. A compound according to any one of the preceding claims wherein R⁵ is selected from hydrogen, fluorine, chlorine, bromine, methyl, ethyl, hydroxyethyl, methoxymethyl, cyano, CF₃, NH₂, NHCOR^{9b} and NHCONHR^{9b} where R^{9b} is phenyl or benzyl optionally substituted by hydroxy, C₁₋₄ acyloxy, fluorine, chlorine, bromine, trifluoromethyl, cyano, C₁₋₄ hydrocarbyloxy and C₁₋₄ hydrocarbyl optionally substituted by C₁₋₂ alkoxy or hydroxy.
- 45. A compound according to any one of the preceding claims wherein R² and R³ are independently selected from hydrogen, C₁₋₄ hydrocarbyl and C₁₋₄ acyl
 - 46. A compound according to claim 28 wherein R² and R³ are independently selected from hydrogen and methyl.
 - 47. A compound according to claim 29 wherein R² and R³ are both hydrogen.
- 48. A compound according to any one of the preceding claims having a molecular weight of less than 525.
- 49. A compound according to claim 1 of the formula (I) which is:
 2-phenyl-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
 3-phenyl-2-[3-(1H-pyrazol-4-yl)-phenyl]-propionitrile;
 2-[4-(3,5-dimethyl-1H-pyrazol-4-yl)-phenyl]-2-phenyl-ethylamine;
 2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
 2-[3-(3,5-dimethyl-1H-pyrazol-4-yl)-phenyl]-1-phenyl-ethylamine;

```
3-phenyl-2-[3-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             3-phenyl-2-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             {3-(4-chloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methyl-amine;
             {3-(3,4-difluoro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methyl-amine;
 5
             {3-(3-chloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methyl-amine;
             3-(4-chloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propionamide;
             3-(4-chloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             3-(3,4-dichloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             4-(4-chloro-phenyl)-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
10
             4-(4-methoxy-phenyl)-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
             4-(4-chloro-phenyl)-1-methyl-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
             4-phenyl-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
             4-[4-(3,5-dimethyl-1H-pyrazol-4-yl)-phenyl]-4-phenyl-piperidine;
             dimethyl-{3-[4-(1H-pyrazol-4-yl)-phenyl]-3-pyridin-2-yl-propyl}-amine;
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-dimethyl-amine;
15
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine (R);
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine (S);
             4-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-morpholine;
20
             4-{4-[1-(4-chloro-phenyl)-2-pyrrolidin-1-yl-ethyl]-phenyl}-1H-pyrazole;
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-isopropyl-amine;
             dimethyl-{2-phenyl-2-[4-(1H-pyrazol-4-yl)-phenyl}-amine;
             {2,2-bis-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-dimethyl-amine;
             {2,2-bis-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
             2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine (R);
25
             2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine (S);
             2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-acetamide;
             1-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-piperazine;
             1-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-piperidine;
             4-{4-[2-azetidin-1-yl-1-(4-chloro-phenyl)-ethyl]-phenyl}-1H-pyrazole;
30
```

```
1-phenyl-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
             2-(4-chloro-phenyl)-N-methyl-2-[4-(1H-pyrazol-4-yl)-phenyl]-acetamide;
             N-methyl-2,2-bis-[4-(1H-pyrazol-4-yl)-phenyl]-acetamide;
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
 5
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-ethyl-amine;
             4-{4-[1-(4-chloro-phenyl)-2-imidazol-1-yl-ethyl]-phenyl}-1H-pyrazole;
             methyl-{2-(4-phenoxy-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-amine;
             {2-(4-methoxy-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
             methyl-{2-[4-(pyrazin-2-yloxy)-phenyl]-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-
10
             amine;
             methyl-{2-phenoxy-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-amine;
             2-{(4-chloro-phenyl)-[4-(1H-pyrazol-4-yl)-phenyl]-methoxy}-ethylamine;
             4-{4-[1-(4-chloro-phenyl)-3-pyrrolidin-1-yl-propyl]-phenyl}-1H-pyrazole;
             4-{4-[3-azetidin-1-yl-1-(4-chloro-phenyl)-propyl]-phenyl}-1H-pyrazole;
15
             methyl-{3-naphthalen-2-yl-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-amine;
             {3-(4-fluoro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methyl-amine;
             4-{4-[4-(4-chloro-phenyl)-piperidin-4-yl]-phenyl}-1H-pyrazole-3-carbonitrile;
             3-(4-phenoxy-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             1-{(4-chloro-phenyl)-[4-(1H-pyrazol-4-yl)-phenyl]-methyl}-piperazine;
20
             1-methyl-4-{phenyl-[4-(1H-pyrazol-4-yl)-phenyl]-methyl}-[1,4]diazepane;
             {3-(3-chloro-phenoxy)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methyl-amine;
             methyl-{2-phenyl-2-[6-(1H-pyrazol-4-yl)-pyridin-3-yl]-ethyl}-amine;
             4-{4-[1-(4-chloro-phenyl)-3-imidazol-1-yl-propyl]-phenyl}-1H-pyrazole;
             4-[4-(3-imidazol-1-yl-1-phenoxy-propyl)-phenyl]-1H-pyrazole;
25
             4-{4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidin-4-yl}-phenol;
             1-{(4-chloro-phenyl)-[4-(1H-pyrazol-4-yl)-phenyl]-methyl}-piperazine;
             {2-(4-fluoro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
             {2-(3-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
             4-[4-(2-methoxy-ethoxy)-phenyl]-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
30
             4-[4-(3-methoxy-propoxy)-phenyl]-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
```

```
3-(3,4-dichloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propionamide;
             2-(4-{2-methylamino-1-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-phenoxy)-
             isonicotinamide;
             {2-(3-chloro-phenoxy)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
 5
             3-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamino}-propan-l-ol;
             2-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamino}-ethanol;
             3-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamino}-propan-1-ol;
             2-{2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamino}-ethanol;
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-cyclopropylmethyl-
10
             amine;
             methyl-[2-[4-(1H-pyrazol-4-yl)-phenyl]-2-(4-pyridin-3-yl-phenyl)-ethyl]-amine;
             4-{3-methylamino-1-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-phenol;
             3-(4-methoxy-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             4-(4-chloro-phenyl)-4-[4-(3-methyl-1H-pyrazol-4-yl)-phenyl]-piperidine;
15
             2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-morpholine;
             (4-{4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidin-4-yl}-phenoxy)-acetic acid;
             4-{4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidin-4-yl}-benzonitrile;
             {2-(4-chloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methyl-amine;
             1-(4-chloro-phenyl)-2-methylamino-1-[4-(1H-pyrazol-4-yl)-phenyl]-ethanol;
20
             2-amino-1-(4-chloro-phenyl)-1-[4-(1H-pyrazol-4-yl)-phenyl]-ethanol;
             4-(3,4-dichloro-phenyl)-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
             4-(3-chloro-4-methoxy-phenyl)-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
             4-(4-chloro-3-fluoro-phenyl)-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
             4-[4-(1H-pyrazol-4-yl)-phenyl]-1,2,3,4,5,6-hexahydro-[4,4']bipyridinyl;
25
             3-(3-chloro-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
             2-methylamino-1-(4-nitro-phenyl)-1-[4-(1H-pyrazol-4-yl)-phenyl]-ethanol;
             2-(3-chloro-4-methoxy-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
             2-(4-chloro-phenyl)-2-fluoro-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
             3-(3,4-dichloro-phenyl)-3-[6-(1H-pyrazol-4-yl)-pyridin-3-yl]-propylamine;
             2-(4-chloro-3-fluoro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
30
```

- 4-(2-chloro-3-fluoro-phenyl)-4-[4-(1H-pyrazol-4-yl)-phenyl]-piperidine;
- 1-{(3,4-dichloro-phenyl)-[4-(1H-pyrazol-4-yl)-phenyl]-methyl}-piperazine;
- 2-(3,4-dichloro-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethylamine;
- {2-(3-chloro-4-methoxy-phenyl)-2-[4-(1H-pyrazol-4-yl)-phenyl]-ethyl}-methyl-amine;
- 4-{4-[2-azetidin-1-yl-1-(4-chloro-phenoxy)-ethyl]-phenyl}-1H-pyrazole;

- 3-(3-chloro-4-methoxy-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propylamine;
- {3-(3-chloro-4-methoxy-phenyl)-3-[4-(1H-pyrazol-4-yl)-phenyl]-propyl}-methylamine;
- 1-{(3,4-dichloro-phenyl)-[4-(1H-pyrazol-4-yl)-phenyl]-methyl}-piperazine; or C-(4-chloro-phenyl)-C-[4-(1H-pyrazol-4-yl)-phenyl]-methylamine; and salts, solvates, tautomers and N-oxides thereof.
 - 50. A compound according to claim 49 which is 2-amino-1-(4-chloro-phenyl)-1-[4-(1H-pyrazol-4-yl)-phenyl]-ethanol or a salt, solvate, tautomer or N-oxide thereof.
- 15 51. A compound according to any one of the preceding claims in the form of a salt, solvate, ester or N-oxide.
 - 52. A compound as defined in any one of claims 1 to 51 for use in medicine.
- 53. A compound as defined in any one of claims 1 to 51 for use in (a) the prophylaxis or treatment of a disease state or condition mediated by protein kinase B; or (b) the prophylaxis or treatment of a disease state or condition mediated by protein kinase A.
 - 54. A compound as defined in any one of claims 1 to 51 for use in the prophylaxis or treatment of a disease state or condition which is selected from a carcinoma of the bladder, breast, colon, kidney, epidermal, liver, lung, oesophagus, gall bladder, ovary, pancreas, stomach, cervix, endometrium, thyroid, prostate, or skin, a hematopoietic tumour of lymphoid lineage, a hematopoietic tumour of myeloid lineage, thyroid follicular cancer, a tumour of mesenchymal origin, a tumour of

the central or peripheral nervous system, melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoctanthoma, thyroid follicular cancer, or Kaposi's sarcoma.

- 55. A compound as defined in any one of claims 1 to 51 for use in the prophylaxis or treatment of a disease state or condition which is selected from breast cancer, ovarian cancer, colon cancer, prostate cancer, oesophageal cancer, squamous cancer and non-small cell lung carcinomas.
 - 56. The use of a compound as defined in any one of claims 1 to 51 for:

10

- (a) the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition mediated by protein kinase B; or
- (b) the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition mediated by protein kinase A; or
- (c) the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition arising from abnormal cell growth;
- (d) the manufacture of a medicament for the prophylaxis or treatment of a disease in which there is a disorder of proliferation, apoptosis or differentiation.
- A pharmaceutical composition comprising a novel compound as defined in any one of claims 1 to 51 and a pharmaceutically acceptable carrier.
- 58. A process for the preparation of a compound of the formula (I) as defined in any one of claims 1 to 51, which process comprises:
 - (a) the reaction of a compound of the formula (X) with a compound of the formula (XI) or an N-protected derivative thereof:

wherein A, E, and R¹ to R⁵ are as defined in any one of claims 1 to 51, one of the groups X and Y is selected from chlorine, bromine, iodine and trifluoromethanesulphonate, and the other one of the groups X and Y is a boronate ester or boronic acid residue, in the presence of a palladium catalyst and a base;

(b) the reductive amination of a compound of the formula (XXXVI):

with HNR²R³ in the presence of a reducing agent; and optionally

(c) the conversion of one compound of the formula (I) into another compound of the formula (I).

European Patent Application No. 04 806 258.2 Astex Therapeutics Limited et al. Our Reference: 40657EP

Basis for amended claims in PCT application as published

Amended claim	Basis in PCT Appl. (PCT claims unless otherwise indicated)	Amended claim	Basis in PCT Appl.
1	Claim 1 + claim 28 (R ⁸) + Claim 42 (R ¹ substituents) + proviso from prior art	30	35
2	Claim 2 + claim 28 (R ⁸) + Claim 42 (R ¹ substituents) + proviso from prior art	31	35
3	3	32	35
4	4	33	36
5	4	34	37 (dependency corrected)
6	. 5	35	38 (dependency corrected)
7	. 6	36	39
8	7	37	. 41
9	. 8	38	40
10	9	39	43
11	10 & 11	40	44
12	12 & 13	41	46
13	14, 15 & 16	42	47
14	17	43	48
15	18	44	49
16	21	45	50
17	22 & 28	46	51
18	23 & 28	47	52

Amended claim	Basis in PCT Appl. (PCT claims unless otherwise indicated)	Amended claim	Basis in PCT Appl.
19 .	24 & 28	48	54
20	26	49	55
21	27 & 28	50	55
22	29	51	56
23	29	52	76
24	. 29	53	57 & 65
25	30	54	Page 92 lines 6 to 21 of description
26	31	55	Page 92 lines 25 to 27 of description
27	32	56	58, 66, 67 & 68
28	33	57	75 (dependency corrected)
29	34	58	77

-

•